

Mass change and motion of a scalar charge in cosmological spacetimes

Roland Haas and Eric Poisson

Department of Physics, University of Guelph, Guelph, Ontario, Canada N1G 2W1

(Dated: November 22, 2004)

Continuing previous work reported in an earlier paper [L.M. Burko, A.I. Harte, and E. Poisson, *Phys. Rev. D* **65**, 124006 (2002)] we calculate the self-force acting on a point scalar charge in a wide class of cosmological spacetimes. The self-force produces two types of effect. The first is a time-changing inertial mass, and this is calculated exactly for a particle at rest relative to the cosmological fluid. We show that for certain cosmological models, the mass decreases and then increases back to its original value. For all other models except de Sitter spacetime, the mass is restored only to a fraction of its original value. For de Sitter spacetime the mass steadily decreases. The second effect is a deviation relative to geodesic motion, and we calculate this for a charge that moves slowly relative to the dust in a matter-dominated cosmology. We show that the net effect of the self-force is to push on the particle. We show that this is not an artifact of the scalar theory: The electromagnetic self-force acting on an electrically charged particle also pushes on the particle. The paper concludes with a demonstration that the pushing effect can also occur in the context of slow-motion electrodynamics in flat spacetime.

I. INTRODUCTION

The prospect of a near-future measurement of gravitational waves produced by the capture of a solar-mass compact object by a massive black hole (one of the identified sources for the Laser Interferometer Space Antenna [1]) is motivating an intense effort to calculate the motion of a compact body in a small-mass-ratio approximation that goes beyond the test-mass description. In this approximation the gravitational influence of the smaller body is taken into account, but the motion is still described as taking place in the background spacetime of the larger body. While the motion is geodesic in the test-mass description, it is accelerated in the small-mass-ratio approximation, and the small body is said to move under the influence of its gravitational self-force. The work that has gone into formulating and evaluating the gravitational self-force is reviewed in detail in Refs. [2, 3], and a shorter, less technical account is given in Ref. [4].

The calculation of the gravitational self-force involves a number of technical and conceptual difficulties, and concrete results pertaining to situations of astrophysical interest have not yet been obtained. To shed light on the physics of self-forces in curved spacetime it is helpful to consider toy problems. In this paper we continue an effort begun in an earlier paper (Ref. [5], hereafter cited as “paper I”) to study the self-force acting on a point scalar charge inserted in a cosmological spacetime. The self-force acting on a scalar charge displays many similarities with the gravitational self-force (including a dependence on the particle’s past history), but it is technically much easier to obtain (fewer field components are involved). And the cosmological models considered here are sufficiently simple that the scalar self-force problem can be solved in a countably infinite number of examples. Beyond these technical advantages, the physics of a scalar point charge is surprisingly interesting: Because a scalar field can radiate in monopole waves, the inertial mass of a scalar charge is not necessarily a constant of the mo-

tion. This phenomenon of mass radiation is one of our main topics in this paper. We also calculate the motion of a scalar charge when it is subjected to its self-force, and compare this with the motion of an electric charge.

The physics of scalar charges in curved spacetime was first investigated by Quinn [6]; his theory is briefly reviewed in Sec. II. In paper I [5] Quinn’s theory was applied to a scalar particle at rest in a spatially-flat cosmological spacetime with metric

$$ds^2 = a^2(\eta)(-d\eta^2 + dx^2 + dy^2 + dz^2), \quad (1.1)$$

where η is conformal time and (x, y, z) are comoving spatial coordinates. In paper I the scale factor was restricted to the forms $a(\eta) = C\eta^2$ (a matter-dominated cosmology) and $a(\eta) = C/|\eta|$ (de Sitter spacetime); C is a scaling constant. For the matter-dominated cosmology it was shown that the scalar charge initially loses a fraction (possibly all) of its mass, but that this radiated mass eventually returns to the particle. For de Sitter spacetime it was shown that the particle radiates all of its mass within a finite proper time.

In this paper we generalize the work reported in paper I [5] on several fronts. We consider a much larger class of cosmological models, those for which $a(\eta) = C|\eta|^\alpha$ with α restricted to (positive and negative) integers. For this class of models we obtain an exact expression for $m(\eta)$, the particle’s mass function. We show that for all values of α except $\alpha = -1$ (de Sitter spacetime), the mass first decreases and then increases back to a fraction of its original value; when α is positive this fraction is in fact equal to unity.

We also calculate the self-force acting on a scalar charge that moves relative to the cosmological fluid. Here our treatment is no longer as general: it is restricted by a slow-motion assumption and to a matter-dominated cosmology. We show that the net effect of the self-force is to push on the particle: When a charged particle and an uncharged particle start with the same initial velocity, the charged particle subsequently moves faster than the

uncharged particle; ultimately both particles are brought to rest by the cosmological expansion. While this conclusion may appear surprising, we show that it is not an artifact of the scalar theory: the same conclusion applies to an electrically charged particle.

We begin in Sec. II with a description of our cosmological spacetimes and a review of Quinn's theory of point scalar charges. The time-changing mass and the motion of the particle can both be determined once the retarded Green's function for the scalar wave equation has been computed; this computation is carried out in Sec. III. In Sec. IV we examine the inertial mass of a scalar charge at rest in our cosmological spacetimes, and establish the behaviors that were described previously. In Sec. V we calculate the motion of a slowly-moving scalar charge in a matter-dominated cosmology, and show that the net effect of the self-force is to push on the particle. In Sec. VI we carry out a similar calculation for an electrically charged particle, but we generalize the cosmological models to the class described by $a(\eta) = C\eta^\alpha$ for any (real or integer, but positive) α ; we show that the electromagnetic self-force also pushes on the particle. In our final section VI we reflect on this perhaps surprising property of the self-force, that it accelerates the particle instead of slowing it down (relative to the motion of an uncharged particle); this goes against a naive expectation that the self-force should always give rise to radiative damping. In Sec. VI we show that this property is not an artifact of curved spacetime: It can be manifested also in the simplest context of slow-motion electrodynamics in flat spacetime.

II. EQUATIONS OF MOTION OF A SCALAR CHARGE IN A COSMOLOGICAL SPACETIME

A particle of mass m and scalar charge q moves in a cosmological spacetime with metric

$$ds^2 = a^2(\eta)(-d\eta^2 + dx^2 + dy^2 + dz^2), \quad (2.1)$$

where η is conformal time and $\mathbf{x} = (x, y, z)$ are comoving spatial coordinates; we use $x \equiv (\eta, \mathbf{x})$ to label a spacetime event, and boldfaced symbols represent three-dimensional vectors in flat space. The cosmology is restricted to be spatially flat, and the scale factor is restricted to the form

$$a(\eta) = C|\eta|^\alpha, \quad (2.2)$$

where C is a scaling constant and α is a constant (positive or negative) exponent. When $\alpha > 0$ we take η to be positive and increasing from 0 to ∞ . When $\alpha < 0$ we take η to be negative and increasing from $-\infty$ to 0. In all cases $a(\eta)$ increases with increasing η . The particle moves on a world line γ described by the parametric relations $z^\mu(\tau)$, with τ denoting proper time; its velocity vector is $u^\mu = dz^\mu/d\tau$.

The mass density of the cosmological fluid is

$$\rho = \frac{3\alpha^2}{8\pi C^2} \frac{1}{|\eta|^{2\alpha+2}} \quad (2.3)$$

and its pressure is given by

$$p = \frac{2-\alpha}{3\alpha}\rho. \quad (2.4)$$

Imposing the dominant energy condition (see, for example, Sec. 9.2 of Ref. [7] or Sec. 2.1 of Ref. [8]) restricts α to the interval $\alpha \geq \frac{1}{2}$ if it is positive, and $\alpha \leq -1$ if it is negative. For $\alpha = 0$ we have flat spacetime, for $\alpha = 1$ a radiation-dominated cosmology ($p = \frac{1}{3}\rho$), for $\alpha = 2$ a matter-dominated cosmology ($p = 0$), and for $\alpha = -1$ we have de Sitter spacetime ($p = -\rho$).

The scalar field Φ produced by the scalar charge q obeys the wave equation

$$g^{\alpha\beta}\nabla_\alpha\nabla_\beta\Phi = -4\pi\mu, \quad (2.5)$$

in which ∇_α indicates covariant differentiation and

$$\mu(x) = q \int_\gamma \delta_4(x, z(\tau)) d\tau \quad (2.6)$$

is the scalar charge density; here $\delta_4(x, x')$ is a scalarized, four-dimensional Dirac distribution that satisfies $\int \delta_4(x, x') \sqrt{-g} d^4x = 1$ when x' is within the domain of integration. The physical solution to Eq. (2.5) is written as

$$\Phi(x) = q \int_\gamma G(x, z(\tau)) d\tau, \quad (2.7)$$

in terms of a retarded Green's function $G(x, x')$ which is a solution to

$$g^{\alpha\beta}\nabla_\alpha\nabla_\beta G(x, x') = -4\pi\delta_4(x, x'). \quad (2.8)$$

When viewed as a function of x' , the Green's function has support both on and within the past light cone of the spacetime event x ; the contribution from the light cone is singular, but the contribution from the interior is smooth.

The field $\Phi(x)$ is singular on γ but its action on the scalar charge is nevertheless well defined. As shown by Quinn [6] (and reviewed in Refs. [2] and [3]), the equations of motion of a freely moving scalar charge are

$$m \frac{Du^\mu}{d\tau} = q^2 (\delta^\mu_\nu + u^\mu u_\nu) \left[\frac{1}{6} R^\nu{}_\lambda u^\lambda + \int_0^{\tau-\epsilon} \nabla^\nu G(z(\tau), z(\tau')) d\tau' \right] \quad (2.9)$$

and

$$\frac{dm}{d\tau} = -q^2 \left[\frac{1}{12} \mathcal{R} + u^\mu \int_0^{\tau-\epsilon} \nabla_\mu G(z(\tau), z(\tau')) d\tau' \right]. \quad (2.10)$$

The left-hand side of Eq. (2.9) is the product of $m(\tau)$ with the particle's covariant acceleration vector, and the right-hand side is the scalar self-force. This involves the spacetime's Ricci tensor and the particle's velocity vector evaluated at the current position $z(\tau)$ on the world line. It also involves the gradient of the retarded Green's function evaluated at $z(\tau)$ and a prior position $z(\tau')$ — the Green's function is differentiated with respect to x before making the identification $x = z(\tau)$. Finally, the self-force involves an integration over the past portion of the particle's world line, from the big bang at $\tau' = 0$ to almost $\tau' = \tau$; the integration is cut short by $\epsilon = 0^+$ to avoid the singular behavior of the Green's function at coincidence. The right-hand side of Eq. (2.10) gives the rate at which the particle's inertial mass changes with proper time τ ; it involves the same objects as Eq. (2.9), together with $\mathcal{R} = R^\mu{}_\mu$, the spacetime's Ricci scalar evaluated at $z(\tau)$.

To solve Eqs. (2.5) and (2.8) in the cosmological spacetimes it is useful to introduce the reduced field

$$\psi(\eta, \mathbf{x}) = a(\eta)\Phi(x), \quad (2.11)$$

the reduced density $\mu^*(\eta, \mathbf{x}) = a^3(\eta)\mu(x)$, and the reduced Green's function

$$g(\eta, \mathbf{x}; \eta', \mathbf{x}') = a(\eta)a(\eta')G(x, x'). \quad (2.12)$$

These satisfy

$$\left(-\frac{\partial^2}{\partial \eta^2} + \nabla^2 + \frac{\alpha(\alpha-1)}{\eta^2}\right)\psi = -4\pi\mu^* \quad (2.13)$$

and

$$\left(-\frac{\partial^2}{\partial \eta^2} + \nabla^2 + \frac{\alpha(\alpha-1)}{\eta^2}\right)g = -4\pi\delta(\eta - \eta')\delta_3(\mathbf{x} - \mathbf{x}'). \quad (2.14)$$

In these equations ∇^2 stands for the Laplacian operator in three-dimensional flat space, and $\delta_3(\mathbf{x} - \mathbf{x}') = \delta(x-x')\delta(y-y')\delta(z-z')$ is an ordinary three-dimensional Dirac distribution. The reduced density obtained from Eq. (2.6) is

$$\mu^*(\eta, \mathbf{x}) = \frac{q}{\gamma}\delta_3(\mathbf{x} - \mathbf{z}(\eta)), \quad (2.15)$$

where $\gamma \equiv a(\eta)d\eta/d\tau = (1 - \mathbf{v} \cdot \mathbf{v})^{-1/2}$, with $\mathbf{v}(\eta) = d\mathbf{z}/d\eta$ denoting the particle's coordinate velocity vector and $\mathbf{v} \cdot \mathbf{v} \equiv (v^x)^2 + (v^y)^2 + (v^z)^2$. The retarded solution to Eq. (2.13) is

$$\psi(\eta, \mathbf{x}) = \int g(\eta, \mathbf{x}; \eta', \mathbf{x}')\mu^*(\eta', \mathbf{x}')d\eta'd^3x', \quad (2.16)$$

and this statement is equivalent to Eq. (2.7).

III. REDUCED GREEN'S FUNCTION

The difficulty in evaluating Eqs. (2.9) and (2.10) resides mostly in the computation of the retarded Green's

function. In paper I [5] the Green's function was obtained for two specific cosmological models, those for which $\alpha = 2$ (matter dominated) and $\alpha = -1$ (de Sitter). In this section we generalize this calculation and compute the Green's function for arbitrary, but integer, values of α .

For convenience we introduce a positive integer l such that

$$\alpha = -l \quad \text{when } \alpha \leq 0 \quad (3.1)$$

and

$$\alpha = l + 1 \quad \text{when } \alpha > 0. \quad (3.2)$$

We notice that with this substitution, $\alpha(\alpha-1) = l(l+1)$. We notice further that a given value of l produces two distinct cosmological models, one with $\alpha_1 = -l \leq 0$ and another with $\alpha_2 = l + 1 > 0$; these models are “equivalent” in the sense that $\alpha_1(\alpha_1-1) = \alpha_2(\alpha_2-1) = l(l+1)$, so that they give rise to the same reduced Green's function. In this sense, flat spacetime ($\alpha = 0$) is equivalent to a radiation-dominated cosmology ($\alpha = 1$), and de Sitter spacetime ($\alpha = -1$) is equivalent to a matter-dominated cosmology ($\alpha = 2$).

It was explained in some detail in paper I [5] that for the cosmological spacetimes considered in this paper, the general structure of the reduced Green's function is

$$g(\eta, \mathbf{x}; \eta', \mathbf{x}') = \frac{\delta(\eta - \eta' - R)}{R} + \theta(\eta - \eta' - R)V(\eta, \mathbf{x}; \eta', \mathbf{x}'), \quad (3.3)$$

where $R \equiv |\mathbf{x} - \mathbf{x}'| \equiv \sqrt{(x-x')^2 + (y-y')^2 + (z-z')^2}$ and $V(\eta, \mathbf{x}; \eta', \mathbf{x}')$ is a smooth two-point function that satisfies the homogeneous form of Eq. (2.14). The first term on the right of Eq. (3.3) is the singular contribution from the light cone; the second term is the contribution from the cone's interior. In flat spacetime $V = 0$, and the support of the Green's function is entirely on the light cone. Because a radiation-dominated universe ($\alpha = 1$) is equivalent to flat spacetime (in the sense defined in the previous paragraph), we have that $V = 0$ in this case as well.¹ In the following we will exclude $l = 0$ from our considerations, and we will compute the reduced Green's function for $\eta - \eta' > R$; in this domain we have $g = V$.

A Fourier-transform method to solve Eq. (2.14) was described in Sec. IV B of paper I [5], and the reader is referred to this paper for details. The core of the method consists of integrating the ordinary differential equation

$$\left(\frac{d^2}{d\eta^2} + k^2 - \frac{l(l+1)}{\eta^2}\right)\hat{g} = 0 \quad (3.4)$$

¹ This conclusion follows also from the fact that $\mathcal{R} = 0$ for a radiation-dominated cosmology, which implies that the scalar wave equation is then conformally invariant. Because the spacetime is conformally related to flat spacetime, the cosmological and Minkowski solutions are related simply by a conformal factor.

for the function $\hat{g}(\eta, \eta'; k)$, subjected to the boundary conditions

$$\hat{g}(\eta = \eta'; k) = 0, \quad \frac{d\hat{g}}{d\eta}(\eta = \eta'; k) = 4\pi. \quad (3.5)$$

The retarded Green's function is then obtained by inverting the Fourier transform, which leads to

$$\begin{aligned} V &= \frac{1}{2\pi^2 R} \int_0^\infty \hat{g}(\eta, \eta'; k) k \sin(kR) dk \\ &= -\frac{1}{2\pi^2 R} \frac{\partial}{\partial R} \int_0^\infty \hat{g}(\eta, \eta'; k) \cos(kR) dk \end{aligned} \quad (3.6)$$

when $\eta - \eta' > R$. The second representation will be more useful for our purposes.

The general solution to Eq. (3.4) is $\hat{g} = Ak\eta j_l(k|\eta|) + Bk\eta n_l(k|\eta|)$, which is expressed in terms of spherical

Bessel functions. The boundary conditions of Eqs. (3.5) determine the constants A and B , and their form can be simplified by invoking the Wronskian-type relations that are satisfied by the Bessel functions (see, for example, Sec. 10.1.6 of Ref. [9]). We find $A = -4\pi|\eta'|n_l(k|\eta'|)$, $B = 4\pi|\eta'|j_l(k|\eta'|)$, and

$$\hat{g} = 4\pi k\eta|\eta'|[j_l(k|\eta'|)n_l(k|\eta|) - n_l(k|\eta'|)j_l(k|\eta|)]. \quad (3.7)$$

After substituting this result into Eq. (3.6) and using the identities (see, for example, Secs. 9.1.2, 10.1.1, and 10.1.12 of Ref. [9]) $j_l(z) = (\pi/2z)^{1/2}J_{l+\frac{1}{2}}(z)$, $n_l(z) = (-1)^{l+1}(\pi/2z)^{1/2}J_{-l-\frac{1}{2}}(z)$, and $\cos(z) = (\pi z/2)^{1/2}J_{-\frac{1}{2}}(z)$, we obtain

$$V = \text{sign}(\alpha)(-1)^l \sqrt{\frac{\pi}{2}} \frac{1}{R} \frac{\partial}{\partial R} \sqrt{\eta\eta'R} \int_0^\infty \sqrt{k} [J_{l+\frac{1}{2}}(k|\eta'|)J_{-l-\frac{1}{2}}(k|\eta|) - J_{-l-\frac{1}{2}}(k|\eta'|)J_{l+\frac{1}{2}}(k|\eta|)] J_{-\frac{1}{2}}(kR) dk, \quad (3.8)$$

in which all functions are Bessel functions of the first kind. The expression also involves $\text{sign}(\alpha) \equiv \text{sign}(\eta) \equiv \eta/|\eta|$; recall that η and η' are both negative when $\alpha < 0$.

The integrals on the right-hand side of Eq. (3.8) can be evaluated in closed form, thanks to Eq. (6.578.1) of Ref. [10]. In the next paragraph we shall describe the calculation of V for $\alpha > 0$ only; a very similar calculation that would return V for $\alpha < 0$ will not be presented.

The first integral on the right-hand side of Eq. (3.8) is proportional to $1/\Gamma(-l)$ and it vanishes when l is an integer. The second integral is equal to

$$-\frac{(\eta/\eta')^l}{\Gamma(-l + \frac{1}{2})\Gamma(l+1)} \sqrt{\frac{2}{\eta\eta'R}} F_4[-l, \frac{1}{2}; -l + \frac{1}{2}, \frac{1}{2}; (\eta'/\eta)^2, (R/\eta)^2],$$

and it is expressed in terms of Appell's hypergeometric function of two variables (see, for example, Sec. 9.18 of Ref. [10]). Because its first argument is a negative integer, the Appell function is in fact a terminating polynomial in the two variables $(\eta'/\eta)^2$ and $(R/\eta)^2$. After cleaning up the Gamma functions we arrive at

$$\begin{aligned} V &= -\frac{(2l-1)!!}{(2l)!!} \frac{(\eta/\eta')^l}{R} \\ &\quad \times \frac{\partial}{\partial R} F_4[-l, \frac{1}{2}; -l + \frac{1}{2}, \frac{1}{2}; (\eta'/\eta)^2, (R/\eta)^2] \end{aligned} \quad (3.9)$$

for the reduced Green's function. There remains the task of evaluating the derivative with respect to R . We use the identity

$$\frac{\partial}{\partial y} F_4(\alpha, \beta; \gamma, \gamma'; x, y) = \frac{\alpha\beta}{\gamma'} F_4(\alpha+1, \beta+1; \gamma, \gamma'+1; x, y),$$

which is easily established from the series representation of the Appell function. Our final result is

$$V(\eta, \mathbf{x}; \eta', \mathbf{x}') = \frac{(2l-1)!!}{(2l-2)!!} \frac{(\eta/\eta')^l}{\eta^2}$$

$$\times F_4[-l+1, \frac{3}{2}; -l + \frac{1}{2}, \frac{3}{2}; (\eta'/\eta)^2, (R/\eta)^2], \quad (3.10)$$

where $R \equiv |\mathbf{x} - \mathbf{x}'|$. The complete reduced Green's function is then given by Eq. (3.3), and $G(x, x')$ can be obtained from Eq. (2.12).

The calculation for $\alpha < 0$ returns a different expression for V , but this can be obtained directly from Eq. (3.10) by performing the exchange $\eta \leftrightarrow \eta'$. These expressions are in fact identical, because the two-point function satisfies the reciprocity relation $V(\eta', \mathbf{x}'; \eta, \mathbf{x}) = V(\eta, \mathbf{x}; \eta', \mathbf{x}')$. This property is derived, for example, in Refs. [2, 3]. We conclude that Eq. (3.10) gives the correct expression for V for both positive and negative values of α .

The Appell function reduces to an ordinary hypergeometric function when the second argument is zero: $F_4(\alpha, \beta; \gamma, \gamma'; x, 0) \equiv F(\alpha, \beta; \gamma; x)$. This implies that Eq. (3.10) simplifies to

$$V(\eta, \mathbf{x}; \eta', \mathbf{x}) = \frac{(2l-1)!!}{(2l-2)!!} \frac{(\eta/\eta')^l}{\eta^2}$$

$$\times F\left[-l+1, \frac{3}{2}; -l+\frac{1}{2}; (\eta'/\eta)^2\right] \quad (3.11)$$

when the spatial points $\mathbf{x}$ and $\mathbf{x}'$ coincide. When the spacetime events x and x' fully coincide we have

$$V(\eta, \mathbf{x}; \eta, \mathbf{x}) = \frac{l(l+1)}{2\eta^2}, \quad (3.12)$$

which is recovered after using identity (15.1.20) of Ref. [9]. This result, when combined with Eqs. (2.12) and (2.2), implies that the full retarded Green's function has the coincidence limit

$$G(\eta, \mathbf{x}; \eta - \epsilon, \mathbf{x}) = \frac{\alpha(\alpha-1)}{2C^2} \frac{1}{\eta^{2\alpha+2}} = \frac{1}{12}\mathcal{R}, \quad (3.13)$$

where $\mathcal{R}$ is the spacetime's Ricci tensor at $x = (\eta, \mathbf{x})$. The presence of $\epsilon = 0^+$ in Eq. (3.13) serves to remind us that the coincidence limit excludes the singular, light-cone contribution to the Green's function; this ϵ plays the same role in Eqs. (2.9) and (2.10).

The general expression of Eq. (3.10) reduces to simple forms when l is a small integer. We list a few in Table I. The result for $l = 1$ was first derived in paper I [5]. The general expression of Eq. (3.10), and the additional special cases listed in the Table, are new.

IV. MASS CHANGE OF A CHARGE AT REST

In this section we calculate the time-changing inertial mass of a scalar charge that is comoving with the cosmological fluid. Such a particle moves on a geodesic of the spacetime, and as we shall see in Sec. V, the right-hand side of Eq. (2.9) vanishes in this case. The right-hand side of Eq. (2.10) does not, however, and we shall find that the particle's mass changes with conformal time η . Without loss of generality we place the scalar charge at the spatial position $\mathbf{x} = \mathbf{0}$.

We begin by deriving a convenient expression for the mass function. Equation (2.10) can be written as

$$\frac{dm}{d\tau} = -q^2 \left[\frac{1}{12}\mathcal{R} + \int_0^{\tau-\epsilon} \frac{d}{d\tau} G(\tau, \tau') d\tau' \right],$$

or as

$$\frac{dm}{d\tau} = -q^2 \left[\frac{1}{12}\mathcal{R} - G(\tau, \tau-\epsilon) \right] - q^2 \frac{d}{d\tau} \int_0^{\tau-\epsilon} G(\tau, \tau') d\tau'.$$

Taking into account Eq. (3.13) we find that the first term vanishes, and that the equation can be directly integrated. This gives

$$m(\tau) = m(\tau_0) - q^2 \int_{\tau_0}^{\tau-\epsilon} G(\tau, \tau') d\tau'. \quad (4.1)$$

We have truncated the domain of integration from $\tau' = 0$ to $\tau' = \tau_0 > 0$ to avoid problems associated with the big bang singularity; in effect we are assuming that the scalar

charge was created at $\tau = \tau_0$ with an initial mass $m(\tau_0)$. As a final step we change the variable of integration from τ' to η' , using the differential relation $d\tau' = a(\eta') d\eta'$ appropriate for a particle at rest, and we re-introduce the reduced Green's function of Eq. (2.12). Taking into account Eq. (3.3) we arrive at a mass function given by

$$m(\eta) = m_0 - \frac{q^2}{a(\eta)} \int_{\eta_0}^{\eta} V(\eta, \mathbf{0}; \eta', \mathbf{0}) d\eta', \quad (4.2)$$

where $m_0 \equiv m(\eta_0)$ is the particle's mass at the moment of creation.

A more concrete expression for $m(\eta)$ is obtained by substituting Eq. (3.11) into Eq. (4.2). The resulting expression simplifies when we change the variable of integration from η' to $\mu \equiv (\eta'/\eta)^2$, which we recognize as the argument of the hypergeometric function. Simple manipulations yield

$$\begin{aligned} \frac{m(\eta)}{m_0} &= 1 - \frac{(2l-1)!!}{(2l-2)!!} \zeta(\eta_0/\eta)^{\alpha+1} \text{sign}(\alpha) \\ &\times \int_{(\eta_0/\eta)^2}^1 \mu^{-(l+1)/2} F\left(-l+1, \frac{3}{2}; -l+\frac{1}{2}; \mu\right) d\mu, \end{aligned} \quad (4.3)$$

where

$$\zeta \equiv \frac{q^2}{2m_0 C |\eta_0|^{\alpha+1}} \quad (4.4)$$

is a dimensionless parameter that determines the strength of the scalar field's contribution to the mass. We recall from Eqs. (3.1) and (3.2) that $\alpha = -l$ when α is negative, and $\alpha = l+1$ when α is positive. We recall also that when $\alpha < 0$, both η and η_0 are negative, which implies that $(\eta_0/\eta)^2 \geq 1$; when $\alpha > 0$ we have instead $(\eta_0/\eta)^2 \leq 1$.

Because the hypergeometric function that appears in Eq. (4.3) is a simple polynomial of degree $l-1$, closed-form expressions for the mass function can easily be obtained for small values of α . We list a few examples in Table II, and Figs. 1 and 2 provide graphs of the selected mass functions.

The plots of Fig. 1 suggest that when α is a positive integer, the mass function decreases at early times, but then increases until it is restored to its original value. Indeed, it is not difficult to show that the property $m(\eta)/m_0 \rightarrow 1$ as $\eta/\eta_0 \rightarrow \infty$ follows directly from Eq. (4.3) when $\alpha > 0$. The plots of Fig. 2, on the other hand, suggest that when $\alpha \leq -2$, only a fraction of the initial mass is eventually returned to the particle. This is confirmed by a similar calculation, which reveals that when $\alpha < 0$,

$$\frac{m(\eta)}{m_0} \rightarrow 1 - \frac{2}{l-1} \frac{(2l-1)!!}{(2l-2)!!} \zeta \quad (4.5)$$

as $\eta/\eta_0 \rightarrow 0$ (recall that $l = -\alpha$). When $\alpha = -1$ (de Sitter spacetime) the mass steadily decreases, and the limit goes to $-\infty$.

TABLE I: Reduced Green's function $V(\eta, \mathbf{x}; \eta', \mathbf{x}')$ for selected values of l , expressed in terms of $R = |\mathbf{x} - \mathbf{x}'|$. These expressions are obtained from Eq. (3.10).

l	α	$V =$
0	$\{0, 1\}$	0
1	$\{-1, 2\}$	$(\eta\eta')^{-1}$
2	$\{-2, 3\}$	$\frac{3}{2}(\eta^2 + \eta'^2 - R^2)(\eta\eta')^{-2}$
3	$\{-3, 4\}$	$\frac{3}{8}[5\eta^4 + 6\eta^2\eta'^2 + 5\eta'^4 - 10(\eta^2 + \eta'^2)R^2 + 5R^4](\eta\eta')^{-3}$
4	$\{-4, 5\}$	$\frac{5}{16}(\eta^2 + \eta'^2 - R^2)[7\eta^4 + 2\eta^2\eta'^2 + 7\eta'^4 - 14(\eta^2 + \eta'^2)R^2 + 7R^4](\eta\eta')^{-4}$




FIG. 1: Mass function for selected positive values of α , plotted as a function of proper time $\tau/\tau_0 = (\eta/\eta_0)^{\alpha+1}$. The mass initially decreases but it returns to its initial value as $\tau \rightarrow \infty$. The graph was obtained from Eq. (4.3) by setting $\zeta = 0.1$.




FIG. 2: Mass function for selected negative values of α , plotted as a function of proper time τ/τ_0 . When $\alpha \neq -1$, $\tau/\tau_0 = (\eta/\eta_0)^{\alpha+1}$, and the mass initially decreases but returns to a fraction of its initial value as $\tau \rightarrow \infty$. When $\alpha = -1$ (de Sitter spacetime), $\tau/\tau_0 = 1 - \ln(\eta/\eta_0)$, and the mass decreases monotonically. The graph was obtained from Eq. (4.3) by setting $\zeta = 0.1$.

TABLE II: Mass function for selected values of α , expressed in terms of $t \equiv \eta/\eta_0$. When $\alpha > 0$ t increases from 1 to ∞ . When $\alpha < 0$ t decreases from 1 to 0. These expressions are obtained from Eq. (4.3).

α	$m(\eta)/m_0 = 1 - \zeta(\dots)$
2	$2 \ln(t)t^{-3}$
3	$3t^{-3} - 3t^{-5}$
4	$\frac{15}{8}t^{-3} + \frac{9}{2} \ln(t)t^{-5} - \frac{15}{8}t^{-7}$
5	$\frac{35}{24}t^{-3} + \frac{45}{8}t^{-5} - \frac{45}{8}t^{-7} - \frac{35}{24}t^{-9}$
-1	$-2 \ln(t)$
-2	$3 - 3t^2$
-3	$\frac{15}{8} - \frac{9}{2} \ln(t)t^2 - \frac{15}{8}t^4$
-4	$\frac{35}{24} + \frac{45}{8}t^2 - \frac{45}{8}t^4 - \frac{35}{24}t^6$

The classical theory of point scalar charges allows the particle's inertial mass to become negative. This may happen when the parameters α , q , m_0 , and η_0 are suitably set ($|\alpha|$ or ζ must be sufficiently large). A more complete theory would provide a description of what happens when m goes to zero. One possibility is that the particle and its scalar charge simply cease to exist when the mass is completely radiated away; this scenario is possible because the theory does not demand scalar-charge conservation. Another possibility is that the particle continues to exist as a massless charged object [11]. We shall remain neutral on the issue of continuing the life of the scalar charge beyond $m = 0$.

V. SELF-FORCE ON A SLOWLY MOVING SCALAR CHARGE

In this section we allow the scalar charge to move slowly relative to the cosmological fluid, we calculate the self-force that is acting on the particle, and we integrate the equations of motion to determine the particle's world line. To keep technicalities at a minimum we adopt a matter-dominated cosmology and choose $\alpha = 2$, and we impose the condition $|\mathbf{v}| \ll 1$, where $\mathbf{v} = d\mathbf{z}/d\eta$ is the particle's coordinate velocity vector. The metric of the spacetime is given by Eq. (2.1) with $a(\eta) = C\eta^2$.

The slow-motion assumption implies that the particle's velocity vector can be approximated by $u^\mu = (1, \mathbf{v})/a(\eta)$.

The spatial components of the covariant acceleration are then

$$\frac{Du^a}{d\tau} = \frac{1}{a^2} \left(\frac{dv^a}{d\eta} + \frac{2}{\eta} v^a \right). \quad (5.1)$$

The velocity of a particle moving on a geodesic therefore satisfies $d\mathbf{v}/d\eta = -2\mathbf{v}/\eta$, and in time the particle is driven toward the same state of motion ($\mathbf{v} = \mathbf{0}$) as the cosmological fluid. From a Newtonian point of view we might say that the cosmological expansion exerts a dragging force on the particle which compels it to come to an eventual rest. We shall adopt this point of view and refer to the $2\mathbf{v}/\eta$ term in the equations of motion as the “cosmological dragging force”.

The self-force acting on a scalar charge can be decomposed into “Ricci” and “tail” forces. The Ricci force appears as the first term on the right-hand side of Eq. (2.9), while the tail force involves an integral over the past portion of the world line. For a matter-dominated cosmology the Ricci tensor is given by $R^x_x = R^y_y = R^z_z = -R^\eta_\eta = 6/(C^2\eta^6)$, and the spatial components of the Ricci force are

$$F_{\text{Ricci}}^a = \frac{1}{a^2} \frac{2q^2}{C\eta^4} v^a. \quad (5.2)$$

Notice that the Ricci force is directed in the direction of $\mathbf{v}$. For a matter-dominated cosmology the retarded Green’s function is given by $G = V/[a(\eta)a(\eta')]$ with $V = 1/(\eta\eta')$, so that $G = C^{-2}(\eta\eta')^{-3}$. Evaluation of the tail integral on the right-hand side of Eq. (2.9) gives

$$F_{\text{tail}}^a = \frac{1}{a^2} \left(-\frac{3q^2}{C\eta^4} \ln \frac{\eta}{\eta_0} \right) v^a \quad (5.3)$$

for the tail force. Notice that we have again truncated the tail integral so that it begins at $\eta = \eta_0$. Notice also that the tail force is directed in the direction opposite to $\mathbf{v}$.

Combining Eqs. (5.1)–(5.3) gives us the particle’s equations of motion,

$$\frac{d\mathbf{v}}{d\eta} = -\frac{2}{\eta} \mathbf{v} + \frac{q^2}{C\eta^4 m(\eta)} \left(2 - 3 \ln \frac{\eta}{\eta_0} \right) \mathbf{v}. \quad (5.4)$$

The first term on the right-hand side of Eq. (5.4) is the cosmological dragging force, which is directed along $-\mathbf{v}$. The second term is the Ricci force, and this is directed along $\mathbf{v}$. Finally, the third term is the tail force, and this is directed along $-\mathbf{v}$. The sign of the self-force depends on time: The Ricci force dominates at early times (η/η_0 close to unity) and it pushes on the particle, while the tail force dominates at late times and pulls on the particle.

We shall assume that q^2 is small and that the self-force produces only a small correction to the particle’s motion. In this regime we can replace $m(\eta)$ by m_0 on the right-hand side of Eq. (5.4), and our precise requirement is that

$$\zeta \equiv \frac{q^2}{2m_0 C\eta_0^3} \ll 1; \quad (5.5)$$

notice that unlike q^2 , ζ is dimensionless. Introducing $t \equiv \eta/\eta_0 \geq 1$ we rewrite Eq. (5.4) as

$$\frac{d\mathbf{v}}{dt} = -\frac{2}{t} \mathbf{v} + \frac{2\zeta}{t^4} (2 - 3 \ln t) \mathbf{v}. \quad (5.6)$$

The largest term on the right of Eq. (5.6) is the cosmological dragging force, and this is only slightly offset by the self-force. Integration of Eq. (5.6), in the regime $\zeta \ll 1$, produces

$$\mathbf{v}(t) = \frac{\mathbf{v}(1)}{t^2} \left[1 + \frac{2\zeta}{3t^3} (t^3 - 1 + 3 \ln t) + O(\zeta^2) \right]. \quad (5.7)$$

This result shows that when a charged ($\zeta \neq 0$) particle and an uncharged ($\zeta = 0$) particle start with the same initial velocity $\mathbf{v}(1)$, the charged particle moves faster at times $t > 1$. Ultimately, however, both particles are brought to rest by the cosmological expansion.

VI. SELF-FORCE ON A SLOWLY MOVING ELECTRIC CHARGE

That a charged particle moves faster than an uncharged particle in a cosmological spacetime (starting with equal initial velocities) is not an artifact of the scalar theory. The same conclusion applies in the context of electrodynamics. In this section we consider a point particle of (constant) mass m and electric charge e that moves slowly relative to the cosmological fluid. We once more calculate the self-force and integrate the particle’s equations of motion. Here we are able to select a more general class of cosmological models, and we let the scale factor take its general form $a(\eta) = C\eta^\alpha$ with α no longer restricted to integers; we do, however, restrict α to be positive in order to keep the discussion simple. We once more impose the condition $|\mathbf{v}| \ll 1$.

It is known that the curved-spacetime version of Maxwell’s theory is conformally invariant (see, for example, Appendix D of Ref. [7]), and the metric of Eq. (2.1) is clearly conformally related to flat spacetime. This implies that the electromagnetic field produced by the moving charge propagates without tails, and that the self-force is purely local. The charge’s equations of motion were first determined by Hobbs [12], who corrected the earlier pioneering treatment of DeWitt and Brehme [13] (see also the reviews [2] and [3] for self-contained derivations). In a conformally-flat spacetime, and in the absence of an external force, the particle moves according to

$$m \frac{Du^\mu}{d\tau} = \frac{1}{3} e^2 (\delta^\mu_\nu + u^\mu u_\nu) R^\nu_\lambda u^\lambda; \quad (6.1)$$

here the self-force contains only a Ricci term, and the tail force is absent.

The nonvanishing components of the Ricci tensor are $R^\eta_\eta = -3\alpha C^{-2}\eta^{-2\alpha-2}$ and $R^x_x = R^y_y = R^z_z = \alpha(2\alpha - 1)C^{-2}\eta^{-2\alpha-2}$. In the slow-motion approximation

the particle's velocity vector is $u^\mu = (1, \mathbf{v})/a(\eta)$, where $\mathbf{v} = d\mathbf{z}/d\eta$ is the coordinate velocity. Using this information, we find that the spatial components of Eq. (6.1) reduce to

$$\frac{d\mathbf{v}}{d\eta} = -\frac{\alpha}{\eta}\mathbf{v} + \frac{2}{3}\alpha(\alpha+1)\frac{e^2}{mC}\frac{1}{\eta^{\alpha+2}}\mathbf{v}. \quad (6.2)$$

The first term on the right-hand side of Eq. (6.2) is the cosmological dragging force; it is directed along $-\mathbf{v}$. The second term is the electromagnetic self-force; it is directed along $\mathbf{v}$ and it pushes on the particle.

In order to integrate Eq. (6.2) we rewrite it as

$$\frac{d\mathbf{v}}{dt} = -\frac{\alpha}{t}\mathbf{v} + \frac{4}{3}\alpha(\alpha+1)\zeta\frac{1}{t^{\alpha+2}}\mathbf{v}, \quad (6.3)$$

re-introducing $t \equiv \eta/\eta_0$ and defining the small dimensionless parameter

$$\zeta = \frac{e^2}{2mC\eta_0^{\alpha+1}}. \quad (6.4)$$

Integration of Eq. (6.3) for $\zeta \ll 1$ is straightforward, and we obtain

$$\mathbf{v}(t) = \frac{\mathbf{v}(1)}{t^\alpha} \left[1 + \frac{4\alpha\zeta}{3t^{\alpha+1}} (t^{\alpha+1} - 1) + O(\zeta^2) \right]. \quad (6.5)$$

This result leads to the same conclusion as in Sec. V: When a charged ($\zeta \neq 0$) particle and an uncharged ($\zeta = 0$) particle start with the same initial velocity $\mathbf{v}(1)$, the charged particle moves faster at times $t > 1$. Both particles are eventually brought to rest by the cosmological expansion.

VII. DISCUSSION: A PUSHING SELF-FORCE?

It is somewhat surprising that a self-force (scalar or electromagnetic) would act so as to push the particle; this goes against a naive expectation that the self-force should always give rise to radiative damping. The examples worked out in the preceding sections illustrate that this expectation is indeed naive. But there is no need to resort to curved-spacetime examples to show this. The expectation is naive also in the simplest context of slow-motion electrodynamics in flat spacetime.

Consider a particle of mass m and electric charge e moving in the field of a charge $-Q$; we take e and Q to be positive quantities, and we suppose that Q is held fixed at the origin of a Cartesian coordinate system. The external force acting on e is the Coulomb force exerted by Q :

$$\mathbf{F}_{\text{ext}} = -k\frac{eQ}{r^2}\hat{\mathbf{r}}, \quad (7.1)$$

where $k = (4\pi\epsilon_0)^{-1}$, $r^2 = x^2 + y^2 + z^2$ is the squared distance from Q , and $\hat{\mathbf{r}} = \mathbf{x}/r$ is a radial unit vector; the

force of Eq. (7.1) is attractive. The self-force acting on e is

$$\mathbf{F}_{\text{self}} = \frac{\mu_0}{6\pi c} \frac{e^2}{m} \dot{\mathbf{F}}_{\text{ext}}, \quad (7.2)$$

where an overdot indicates differentiation with respect to time. This expression for the self-force is derived, for example, in Sec. 75 of the Landau and Lifshitz textbook [14]. It is usually expressed in terms of the rate of change of the particle's acceleration vector, but Landau and Lifshitz explain that it is consistent to make the replacement $\mathbf{a} \rightarrow \mathbf{F}_{\text{ext}}/m$ on the right-hand side of Eq. (7.2). A first-principles derivation of this equation is also presented in Ref. [15].

From Eq. (7.1) we calculate that the rate of change of the external force is given by $\dot{\mathbf{F}}_{\text{ext}} = -keQ(\mathbf{v} - 3\dot{r}\hat{\mathbf{r}})/r^3$, where $\dot{r} \equiv \mathbf{v} \cdot \hat{\mathbf{r}}$ is the radial component of the velocity vector. We decompose this vector in terms of radial and tangential components,

$$\mathbf{v} = \mathbf{v}_\perp + \dot{r}\hat{\mathbf{r}}, \quad \mathbf{v}_\perp \cdot \hat{\mathbf{r}} = 0, \quad (7.3)$$

and we substitute this result into Eq. (7.2). This gives

$$\mathbf{F}_{\text{self}} = \frac{\mu_0 k}{6\pi c} \frac{e^3 Q}{m} (-\mathbf{v}_\perp + 2\dot{r}\hat{\mathbf{r}}) \quad (7.4)$$

for the self-force acting on the moving charge.

Suppose that the charge e is moving tangentially, so that $\dot{r} = 0$. Then the self-force is directed along $-\mathbf{v}_\perp = -\mathbf{v}$ and we find that it is indeed a dragging force. Now suppose that the charge is moving radially inward, so that $\mathbf{v}_\perp = \mathbf{0}$ and $\dot{r} < 0$. Then the self-force is directed along $-\hat{\mathbf{r}}$, that is, in the same direction as the external force. In this situation the self-force pushes on the particle, in violation of the naive expectation.

More generally, the rate of work done by the self-force, $\mathbf{F}_{\text{self}} \cdot \mathbf{v}$, is proportional to $-\mathbf{v}_\perp^2 + 2\dot{r}^2$, where $\mathbf{v}_\perp \equiv |\mathbf{v}_\perp|$. When $\mathbf{v}_\perp > \sqrt{2}\dot{r}$ the work done on the particle is negative and the self-force exerts a drag. When, however, $\mathbf{v}_\perp < \sqrt{2}\dot{r}$ the work done is positive and the self-force pushes.

Does a pushing self-force violate energy conservation? The answer is in the negative, in spite of appearances to the contrary. Quite generally, the rate of work done on a particle of charge e by the self-force (which we now express in terms of the particle's acceleration $\mathbf{a}$) is

$$\mathbf{F}_{\text{self}} \cdot \mathbf{v} = \frac{\mu_0 e^2}{6\pi c} \dot{\mathbf{a}} \cdot \mathbf{v} = \frac{\mu_0 e^2}{6\pi c} \left[\frac{d}{dt}(\mathbf{a} \cdot \mathbf{v}) - |\mathbf{a}|^2 \right].$$

The last term on the right-hand side gives the electromagnetic power radiated by the particle, and we have the statement

$$[\text{rate of work done}] + [\text{radiated power}] = \frac{\mu_0 e^2}{6\pi c} \frac{d}{dt}(\mathbf{a} \cdot \mathbf{v}). \quad (7.5)$$

Integrating this equation over a finite time interval gives rise to the following conservation statement:

$$[\text{work done}] + [\text{radiated energy}] = [\text{total change in } \mu_0 e^2(\mathbf{a} \cdot \mathbf{v})/(6\pi c)]. \quad (7.6)$$

The term on the right can be interpreted as a change in the system's "internal energy" as it moves from one configuration to another. (The system is the charged particle plus the agent responsible for the external force.) This form of energy is associated with the part of the electromagnetic field that stays bound to the system instead of propagating freely as electromagnetic radiation.

In some circumstances the total change in internal energy is zero. For example, when the motion is periodic the total change vanishes after one complete period. As another example, the total change vanishes when the motion proceeds from a state of no acceleration to another state of no acceleration. In such circumstances the statement of Eq. (7.6) simplifies to

[work done] + [radiated energy] = 0, and the work done by the self-force must be negative. In more general circumstances, however, the statement of energy conservation must include the change in internal energy, and it becomes possible for the self-force to do positive work on the particle. This was well illustrated in the previous discussion.

Acknowledgments

This work was supported by the Natural Sciences and Engineering Research Council of Canada.

-
- [1] The LISA web site is located at <http://lisa.jpl.nasa.gov/>.
 - [2] E. Poisson, *The motion of point particles in curved spacetime*, Living Rev. Relativity **7** (2004), 6. [Online article]: cited on February 7, 2008, <http://www.livingreviews.org/lrr-2004-6>.
 - [3] E. Poisson, *Radiation reaction of point particles in curved spacetime*, Class. Quantum Grav. **21**, R153 (2004).
 - [4] E. Poisson, *The gravitational self-force* (2004), <http://xxx.lanl.gov/abs/gr-qc/0410127>.
 - [5] L. M. Burko, A. I. Harte, and E. Poisson, *Mass loss by a scalar charge in an expanding universe*, Phys. Rev. D **65**, 124006 (2002).
 - [6] T. C. Quinn, *Axiomatic approach to radiation reaction of scalar point particles in curved spacetime*, Phys. Rev. D **62**, 064029 (2000).
 - [7] R. W. Wald, *General Relativity* (Chicago University Press, Chicago, 1984).
 - [8] E. Poisson, *A relativist's toolkit: The mathematics of black-hole mechanics* (Cambridge University Press, Cambridge, England, 2004).
 - [9] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover, New York, 1972).
 - [10] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products* (Academic Press, Orlando, 1980).
 - [11] P. Kazinski and A. Sharapov, *Radiation reaction for a massless charged particle*, Class. Quantum Grav. **20**, 2715 (2003).
 - [12] J. M. Hobbs, *A vierbein formalism for radiation damping*, Ann. Phys. (N.Y.) **47**, 141 (1968).
 - [13] B. S. DeWitt and R. W. Brehme, *Radiation damping in a gravitational field*, Ann. Phys. (N.Y.) **9**, 220 (1960).
 - [14] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields, Fourth Edition* (Butterworth-Heinemann, Oxford, England, 2000).
 - [15] E. Poisson, *An introduction to the Lorentz-Dirac equation* (1999), <http://xxx.lanl.gov/abs/gr-qc/9912045>.